

Help file for `ppml_panel_sg`

by Thomas Zylkin

Title

`ppml_panel_sg` - Fast Poisson Pseudo-Maximum Likelihood (PPML) regression for panel “gravity” models with time-varying origin and destination fixed effects and time-invariant pair fixed effects.

Syntax

```
ppml_panel_sg depvar [indepvars] [if] [in], exporter(exp_id) importer(imp_id) year(time_id) [options]
```

exp_id, *imp_id*, and *time_id* are variables that respectively identify the origin, destination, and time period associated with each observation.

Description

`ppml_panel_sg` enables faster computation of the many fixed effects required for panel PPML structural gravity estimation. In particular, it addresses the large number of “pair-wise” FEs needed to consistently identify the effects of time-varying trade policies such as regional trade agreements (see, e.g., Baier & Bergstrand, 2007; Dai, Yotov, & Zylkin, 2014). It also simultaneously absorbs the origin-by-time and destination-by-time FEs implied by theory.

Some options and features of interest:

1. Programmed to run in Mata, making it much faster than existing Stata Poisson commands for estimating the effects of trade policies.
2. Can store the estimated fixed effects in Stata’s memory (but as a single column each, rather than as a large matrix with many zeroes)
3. In addition to the pair fixed effects, also readily allows for specifications with additional pair-specific linear time trends.
4. Supports multi-way clustering of standard errors.
5. All fixed effects are also allowed to vary by industry, in case you wish to examine industry-level variation in trade flows.
6. Can be used with strictly cross-sectional regressions (i.e., without pair fixed effects).
7. Performs Santos Silva & Tenreyro (2010)’s recommended check for possible non-existence of estimates.

Main Options

<u>nopair</u>	Use origin-time and destination-time fixed effects only (do not include pair fixed effects).
<u>trend</u>	Add linear, pair-specific time trends.
<u>symmetric</u>	Assume pair fixed effects apply symmetrically to flows in both directions, as in, e.g., Anderson & Yotov (2016). Time trends, if specified, will also be symmetric.
<u>industry</u> (<i>ind_id</i>)	<i>ind_id</i> is a varname identifying the industry associated with each observation. When an <i>ind_id</i> is provided, the origin-time fixed effects become origin-industry-time effects, the destination-time fixed effects become destination-industry-time effects, and pair-specific terms become origin-destination-industry-specific.
<u>offset</u> (<i>varname</i>)	Include <i>varname</i> as a regressor with coefficient constrained to be equal to 1.
<u>cluster</u> (<i>cluster_id</i>)	Specifies clustered standard errors, clustered by <i>cluster_id</i> . The default is clustering by <i>exp_id-imp_id</i> , unless nopair is enabled. For multi-way clustering, <i>cluster_id</i> may be a varlist (max length is 4.)
<u>robust</u>	Use robust standard errors. This is the default if nopair is enabled.
<u>dropsingletons</u>	Drop “singletons” beforehand à la reghdfe .
<u>multiway</u>	Automatic three-way clustering by <i>exp-id</i> , <i>imp-id</i> , and <i>time-id</i> .
<u>nosterr</u>	Do not compute standard errors (saves time if you only care about point estimates).
<u>tolerance</u> (#)	The default tolerance is 1e-12.
<u>maxiter</u> (#)	The default maximum number of iterations is 10,000.
<u>noaccel</u>	Do not use “acceleration” when computing estimates.
<u>verbose</u> (#)	Show iterative output for every #th iteration. Default is 0 (no output).

Guessing and Storing Values

These options allow you to store results for fixed effects in memory as well as use information from memory to set up initial guesses. You may also utilize the “multilateral resistances” of Anderson & van Wincoop (2003).

<u>olsguess</u>	Use reghdfe to initialize guesses for coefficient values.
<u>guessB</u> (<i>str</i>)	Supply the name of a row vector with guesses for coefficient values.
<u>guessD</u> (<i>varname</i>)	Guess initial values for the (exponentiated) set of pair fixed effects. Default is all 1s.
<u>guessS</u> (<i>varname</i>)	Guess initial values for the (exponentiated) set of origin-time fixed effects. Default is the share of depvar within each [<i>ind_id</i>]- <i>time-id</i> associated with each <i>exp-id</i> .

- guessM**(*varname*) Guess initial values for the (exponentiated) set of destination-time fixed effects. Default is the share of `depvar` within each [*ind_id*-]*time-id* associated with each *imp-id*.
- guessO**(*varname*) Guess initial values for the set of “outward” multilateral resistances. Default is all 1s. Overrides **genS**.
- guessI**(*varname*) Guess initial values for the set of “inward” multilateral resistances. Default is all 1s. Overrides **genM**.
- guessTT**(*varname*) Guess initial values for pair time trends. These are not exponentiated. Default is all 0s.
- genD**(*newvar*), **genS**(*newvar*), **genM**(*newvar*), **genO**(*newvar*), **genI**(*newvar*), and **genTT**(*newvar*): These options store fixed effects and/or time trend parameters in memory as new variables. To store predicted values, use **predict**(*newvar*).

Check for Existence

As with `ppml`, `ppml_panel_sg` checks your specification beforehand to ensure that valid estimates will indeed exist. These options affect how this check is performed.

- nocheck** Do not check for existence.
- strict** Applies a more conservative set of exclusion conditions when checking whether each regressor may be included; mimics “strict” option from `ppml`.
- keep** Keeps observations that are perfectly predicted by excluded regressors; mimics “keep” option from `ppml`.

Background

As a typical application, consider the following PPML regression:

$$X_{ijt} = \exp [\ln S_{it} + \ln M_{jt} + \ln D_{ij} + b \times RTA_{ijt}] + e_{ijt}. \quad (1)$$

X_{ijt} are international trade flows. i , j , and t are indices for origin, destination, and time. The goal is to consistently estimate the average effect of RTA_{ijt} , a dummy variable for the presence of a regional trade agreement on trade flows, using a “structural gravity” specification. The origin-time and destination-time fixed effects— S_{it} and M_{jt} —ensure the theoretical restrictions implied by structural gravity are satisfied. The pair fixed effect— D_{ij} —then absorbs all time-invariant pair characteristics that may be correlated with the likelihood of forming an RTA.

Computationally, the biggest obstacle to estimating (1) is the pair fixed effect term D_{ij} . Because a unique D_{ij} must be computed for each pair, the number of D_{ij} ’s increases rapidly with the number of locations. For a balanced international trade data set with 75 countries trading with each other over 10 years (not an especially large sample for trade data), there will be on the order of $75^2 = 5,625$ pair fixed effects that must be computed. In addition (ignoring collinearity), we will also require $75 \times 2 \times 10 = 1,500$ origin-time and destination-time effects. The

total number of parameters needed to estimate (1) (around 7,000) would normally require a long computing time in Stata, likely several hours at least. If we push the number of locations and/or years further, we will quickly approach Stata's matsize limits, beyond which estimation becomes infeasible.

To date, this is the only available Stata command that will perform "fast" estimation of specifications such as (1) using PPML. It works by manipulating the first order conditions of the Poisson to produce analytical expressions for each of the fixed effects that can be computed via simple iteration. In this way, it both adapts and extends existing procedures described in Guimarães & Portugal (2010) and Figueiredo, Guimarães, & Woodward (2015) for estimating Poisson models with high dimensional fixed effects. These works and others are recommended below for further reading.

Examples

To perform a basic panel estimation such as (1):

```
ppml_panel_sg trade rta, ex(iso_o) im(iso_d) y(year)
```

To add pair-specific time trends, i.e.,

$$X_{ijt} = \exp [\ln S_{it} + \ln M_{jt} + \ln D_{ij} + a_{ij} \times t + b \times RTA_{ijt}] + e_{ijt}, \quad (2)$$

you would input:

```
ppml_panel_sg trade rta, ex(iso_o) im(iso_d) y(year) trend
```

If you want your pair fixed effects to be symmetric (i.e., $D_{ij} = D_{ji}$), the syntax is:

```
ppml_panel_sg trade rta, ex(iso_o) im(iso_d) y(year) sym
```

To estimate coefficients of more traditional, time-invariant gravity variables, such as bilateral distance, use the **nopair** option:

```
ppml_panel_sg trade ln_dist colony language contiguity rta,  
ex(iso_o) im(iso_d) y(year) nopair
```

Unlike the regressions with pair fixed effects, however, obtaining estimates for time-invariant regressors may not be noticeably faster than existing methods (e.g., [glm](#), [ppml](#)) unless the number of origin-time and destination-time effects is sufficiently large. You may also exclude the year ID in this last specification if your data includes only 1 year.

Advisory

This estimation command is strictly intended for settings where the dependent variable is spatial flows from one set of locations to another (such as international trade or migration flows). It is not a generalized Poisson fixed effects command. For more general problems that require Poisson estimation, you may try: [poisson](#), [glm](#), [ppml](#),

[xtpoisson](#), [xtpqml](#), and/or [poi2hdfe](#). For an OLS command that can compute similar “gravity” specifications using OLS, I recommend [reghdfe](#).

This is version 1.1 of this command. If you believe you have found an error that can be replicated, or have other suggestions for improvements, please feel free to [contact me](#).

Acknowledgements

I have adapted parts of my code from several other related commands. These include: [poi2hdfe](#), by Paulo Guimarães, [SILS](#), by Keith Head and Thierry Mayer, and [reghdfe](#) by Sergio Correia. I give the utmost credit to each of these authors for creating these programs and making their code available. The inclusion of multi-way clustering in recent versions of `ppml_panel_sg` (from v. 1.1 onward) is thanks to contributions by Joschka Wanner and Mario Larch.

I also thank Yoto Yotov, Davin Chor, Sergio Correia, Paulo Guimarães, and João Santos Silva for kindly taking the time to offer feedback and suggest improvements.

Suggested citation

If you are using this command in your research I would appreciate if you would cite

- Larch, Mario, Wanner, Joschka, Yotov, Yoto V., & Zylkin, Thomas (2017), “The Currency Union Effect: A PPML Re-assessment with High-dimensional Fixed Effects”, Drexel University School of Economics Working Paper 2017-07.

which also provides a technical companion for those interested in understanding how the command works.

Further Reading

- Structural gravity: Anderson & van Wincoop (2003); Head & Mayer (2014)
- On the use of PPML to estimate gravity equations: Santos Silva & Tenreyro (2006, 2011); Arvis & Shepherd (2013); Egger & Staub (2014); Fally (2015)
- Possible non-existence of Poisson MLE estimates: Santos Silva & Tenreyro (2010)
- Consistently estimating the effects of trade policies: Baier & Bergstrand (2007); Dai, Yotov, & Zylkin (2014); Anderson & Yotov (2016); Piermartini & Yotov (2016)
- Estimating models with high dimensional fixed effects: Guimarães & Portugal (2010); Gaure (2011); Figueiredo, Guimarães, & Woodward (2015); Correia (2016)
- Multi-variate Steffenson’s acceleration (used to accelerate convergence): Nievergelt (1991)
- Multi-way clustering: Cameron, Gelbach, & Miller (2011); Egger & Tarlea (2015)
- Singletons: Correia (2015).

References

- Anderson, J. E. & van Wincoop, E. (2003), “Gravity with Gravitas: A Solution to the Border Puzzle”, *American Economic Review* **93**(1), 170–192.
- Anderson, J. E. & Yotov, Y. V. (2016), “Terms of Trade and Global Efficiency Effects of Free Trade Agreements, 1990-2002”, *Journal of International Economics* (**in press**).
- Arvis, J.-F. & Shepherd, B. (2013), “The Poisson quasi-maximum likelihood estimator: a solution to the ‘adding up’ problem in gravity models”, *Applied Economics Letters* **20**(6), 515–519.
- Baier, S. L. & Bergstrand, J. H. (2007), “Do free trade agreements actually increase members’ international trade?”, *Journal of International Economics* **71**(1), 72–95.
- Cameron, A. C., Gelbach, J. B., & Miller, D. L. (2011), “Robust Inference With Multiway Clustering”, *Journal of Business & Economic Statistics* **29**(2), 238–249.
- Correia, S. (2015), “Singletons, Cluster-Robust Standard Errors and Fixed Effects: A Bad Mix”, Mimeo, Duke University.
- Correia, S. (2016), “A Feasible Estimator for Linear Models with Multi-Way Fixed Effects”, Mimeo, Duke University.
- Dai, M., Yotov, Y. V., & Zylkin, T. (2014), “On the trade-diversion effects of free trade agreements”, *Economics Letters* **122**(2), 321–325.
- Egger, P. H. & Staub, K. E. (2014), “GLM estimation of trade gravity models with fixed effects”, Mimeo, ETH Zurich.
- Egger, P. H. & Tarlea, F. (2015), “Multi-way clustering estimation of standard errors in gravity models”, *Economics Letters* **134**, 144–147.
- Fally, T. (2015), “Structural gravity and fixed effects”, *Journal of International Economics* **97**(1), 76–85.
- Figueiredo, O., Guimarães, P., & Woodward, D. (2015), “Industry localization, distance decay, and knowledge spillovers: Following the patent paper trail”, *Journal of Urban Economics* **89**(C), 21–31.
- Gaure, S. (2011), “OLS with Multiple High Dimensional Category Dummies”, Mimeo 14/2010, Oslo University, Department of Economics.
- Guimarães, P. & Portugal, P. (2010), “A simple feasible procedure to fit models with high-dimensional fixed effects”, *Stata Journal* **10**(4), 628–649.
- Head, K. & Mayer, T. (2014), “Gravity equations: Workhorse, toolkit, and cookbook”, *Handbook of International Economics* **4**, 131–196.
- Nievergelt, Y. (1991), “Aitken’s and Steffensen’s accelerations in several variables”, *Numerische Mathematik* **59**(1), 295–310.
- Piermartini, R. & Yotov, Y. (2016), “Estimating Trade Policy Effects with Structural Gravity”, Working Paper ERSD-2016-10, WTO.

- Santos Silva, J. M. C. & Tenreyro, S. (2006), “The Log of Gravity”, *Review of Economics and Statistics* **88**(4), 641–658.
- Santos Silva, J. M. C. & Tenreyro, S. (2010), “On the existence of the maximum likelihood estimates in Poisson regression”, *Economics Letters* **107**(2), 310–312.
- Santos Silva, J. M. C. & Tenreyro, S. (2011), “Further simulation evidence on the performance of the Poisson pseudo-maximum likelihood estimator”, *Economics Letters* **112**(2), 220–222.